arXiv:0910.0566v1 [math.MG] 3 Oct 2009

AN INTERVAL VERSION OF SEPARATION BY SEMISPACES IN MAX-MIN CONVEXITY

VIOREL NITICA AND SERGEĬ SERGEEV

Abstract. In this paper we study separation of a closed box from a max-min convex set by max-min semispaces. This can be regarded as an interval extension of the known separation results. We give a constructive proof of the separation in the case when the box satisfies a certain condition, and we show that the separation is never possible when the condition is not satisfied. We also study the separation of two max-min convex sets by a box and by a box and a semispace. *Key Words:* fuzzy algebra; tropical convexity; interval; separation; *Mathematics Subject Classification 2000:* Primary 52A01; Secondary: 52A30, 08A72

1. Introduction

Consider the set $\mathcal{B} = [0, 1]$ endowed with the operations $\oplus = \max, \wedge = \min$. This is a well-known distributive lattice, and like any distributive lattice it can be considered as a semiring equipped with addition $\oplus$ and multiplication $\otimes := \wedge$. Importantly, both operations are idempotent, $a \oplus a = a$ and $a \otimes a = a \wedge a = a$, and closely related to the order: $a \oplus b = b \Leftrightarrow a \leq b \Leftrightarrow a \wedge b = a$. For standard literature on lattices and semirings see e.g. [2] and [10].

We consider $\mathcal{B}^n$, the cartesian product of n copies of $\mathcal{B}$, and equip this cartesian product with the operations of taking componentwise $\oplus$: $(x \oplus y)_i := x_i \oplus y_i$ for $x, y \in \mathcal{B}^n$ and $i = 1, \ldots, n$, and scalar $\wedge$-multiplication: $(a \wedge x)_i := a \wedge x_i$ for $a \in \mathcal{B}$, $x \in \mathcal{B}^n$ and $i = 1, \ldots, n$. Thus $\mathcal{B}^n$ is considered as a semimodule over $\mathcal{B}$ [10]. Alternatively, one may think in terms of vector lattices [2].

A subset C of $\mathcal{B}^n$ is said to be *max-min convex* if the relations $x, y \in C, \alpha, \beta \in \mathcal{B}, \alpha \oplus \beta = 1$ imply $(\alpha \wedge x) \oplus (\beta \wedge y) \in C$.

This research is supported by NSF grant DMS-0500832 (V. Nitica) as well as EPSRC grant RRAH12809 and RFBR grant 08-01-00601 (S. Sergeev).

The interest in max-min convexity is motivated by the study of tropically convex sets, analogously defined over the semiring $\mathbb{R}_{\max}$, which is the completed set of real numbers $\mathbb{R} \cup \{-\infty\}$ endowed with operations of idempotent addition $a \oplus b := \max(a, b)$ and multiplication $a \otimes b := a + b$. Constructed in [22, 23], tropical convexity and its lattice-theoretic generalizations received much attention and rapidly developed over the last decades [1, 4, 5, 7, 11, 15, 16]. Another source of interest comes from the matrix algebra developed over the max-min semiring, see [3, 9, 19] and references therein.

In this article we continue the study of max-min convex structures started in [17, 18, 13, 14]. We are interested in separation of max-min convex sets by semispaces.

The set

$$\begin{aligned} [x, y]_M &= \{(\alpha \wedge x) \oplus (\beta \wedge y) \in \mathcal{B}^n \,|\, \alpha, \beta \in \mathcal{B}, \alpha \oplus \beta = 1\} = \\ &= \{\max(\min(\alpha, x), \min(\beta, y)) \in \mathcal{B}^n \,|\, \alpha, \beta \in \mathcal{B}, \max(\alpha, \beta) = 1\}, \end{aligned} \tag{1}$$

is fundamental for max-min convexity, it is called the *max-min segment* (or briefly, the *segment*) *joining* x *and* y. As in the ordinary convexity in the real linear space, a set is max-min convex if and only if any two points are contained in it together with the max-min segment joining them. The max-min segments have been described in [17, 20].

Other types of convex sets are max-min semispaces, hemispaces, halfspaces and hyperplanes [18, 13, 14].

For $z \in \mathcal{B}^n$, we call a subset S of $\mathcal{B}^n$ a *max-min semispace* (or, briefly, a *semispace*) *at* z, if it is a maximal (with respect to set-inclusion) max-min convex set avoiding z. Semispaces come from the abstract convexity, see e.g. [21]. One of their main application is in separation results: the family of semispaces is the smallest intersectional basis for the family of all convex sets. We recall that in $\mathcal{B}^n$ there exist at most $n+1$ semispaces at each point, exactly $n+1$ at each finite point, and each convex set avoiding z is contained in at least one of those semispaces [18].

Another object introduced in abstract convexity is the *hemispace*: this is any convex set whose complement is also convex. When the system of convex sets satisfies Pasch axiom, which is the case for max-min convexity [17], a theorem of Kakutani tells us that for any two nonintersecting convex sets C_1 and C_2 there exists a hemispace H containing C_1 such



FIGURE 1. Separation types, $n = 2$

that the complement of H contains C_2. In general, the proof of Kakutani theorem is non-constructive and uses Zorn's Lemma. A constructive proof of this theorem in max-min convexity is, to the authors' knowledge, an open problem.

Hyperplanes and halfspaces are defined by linear forms (in our case, max-min linear). In the max-min case, these sets are in general unrelated to semispaces and they cannot separate a point from a max-min convex set [13, 14]. This is in contrast with very optimistic results in the tropical convexity and its lattice-theoretic generalizations [4, 5, 7, 8, 22], which behave like the ordinary convexity in linear spaces in this respect.

In this paper we study the following interval version of the semispace separation: given a box B, i.e. a Cartesian product of closed intervals, and a max-min convex set C, decide whether it is possible to construct a semispace which contains C and avoids B. In Section 2 we give our main result, Theorem 1, which shows that such separation is indeed possible when B satisfies a certain condition. This condition holds true in particular when B does not contain points with coordinates equal to 1, or when B is reduced to a point. When the condition is not satisfied, we show that the separation by semispaces is never possible. However, separation can be saved if we also allow hemispaces of a certain kind. As a corollary of Theorem 1, we also recover the description of semispaces due to Nitica and Singer [18]. In Section 3 we study the separation of two convex sets by a box and by a box and a semispace. We show that this separation is always possible in $\mathcal{B}^2$, and we provide a counterexample in $\mathcal{B}^3$.

Figure 1 summarizes the types of separation considered in this paper. The convex sets that need to be separated are colored in black, and the separating boxes or semispaces are colored in gray. The sets C_1, C_2 and C are convex and B is a box.

The tropical interval linear algebra was introduced in [12] and [6, Chapter 6]. From this point of view, the present paper may be seen as related to yet undeveloped area of the interval tropical convexity.

2. Separation of boxes from max-min convex sets

For any point $x^0 = (x_1^0, \dots, x_n^0) \in \mathcal{B}^n$ we define a family of sets $S_0(x^0), \dots, S_n(x^0)$ in $\mathcal{B}^n$. The sets are introduced in [18, Proposition 4.1]. Recall that x^0 is called *finite* if it has all coordinates different from zeros and ones. Without loss of generality we may assume that:

$$x_1^0 \geq \cdots \geq x_n^0. \tag{2}$$

The set $\{x_1^0, \dots, x_n^0\}$ admits a natural subdivision into ordered subsets such that the elements of each subset are either equal to each other or are in strictly decreasing order, say

$$\begin{aligned} x_1^0 = \cdots = x_{k_1}^0 > \cdots > x_{k_1+l_1+1}^0 = \cdots = x_{k_1+l_1+k_2}^0 > \cdots \\ > x_{k_1+l_1+k_2+l_2+1}^0 = \cdots = x_{k_1+l_1+k_2+l_2+k_3}^0 > \cdots \\ > x_{k_1+l_1+\cdots+k_{p-1}+l_{p-1}+1}^0 = \cdots = x_{k_1+l_1+\cdots+k_{p-1}+l_{p-1}+k_p}^0 \\ > \cdots > x_{k_1+l_1+\cdots+k_p+l_p}^0 (= x_n^0). \end{aligned} \tag{3}$$

Let us introduce the following notations:

$$L_0 = 0, K_1 = k_1, L_1 = K_1 + l_1 = k_1 + l_1, \tag{4}$$

$$K_j = L_{j-1} + k_j = k_1 + l_1 + ... + k_{j-1} + l_{j-1} + k_j \quad (j = 2, ..., p), \tag{5}$$

$$L_j = K_j + l_j = k_1 + l_1 + ... + k_j + l_j \quad (j = 2, ..., p); \tag{6}$$

we observe that $l_j = 0$ if and only if $K_j = L_j$.

We are ready to define the sets. We need to distinguish the cases when the sequence (3) ends with zeros or begin with ones, since some sets S_i become empty in that case.

Definition 1. *a) If x^0 is finite, then:*

$$S_0(x^0) = \{x \in \mathcal{B}^n | x_i > x_i^0 \text{ for some } 1 \leq i \leq n\}, \tag{7}$$

$$\begin{gathered} S_{K_j+q}(x^0) = \{x \in \mathcal{B}^n | x_{K_j+q} < x^0_{K_j+q}, \text{ or } x_i > x_i^0 \text{ for some } K_j + q + 1 \leq i \leq n\} \\ (q = 1, ..., l_j; j = 1, ..., p) \text{ if } l_j \neq 0, \end{gathered} \tag{8}$$

$$\begin{gathered} S_{L_{j-1}+q}(x^0) = \{x \in \mathcal{B}^n | x_{L_{j-1}+q} < x^0_{L_{j-1}+q}, \text{ or } x_i > x_i^0 \text{ for some } K_j + 1 \leq i \leq n\} \\ (q = 1, ..., k_j; j = 1, ..., p \text{ if } k_1 \neq 0, \text{ or } j = 2, ..., p \text{ if } k_1 = 0). \end{gathered} \tag{9}$$

b) If there exists an index $i \in \{1, ..., n\}$ such that $x_i^0 = 1$, but no index j such that $x_j^0 = 0$, then the sets are $S_1, ..., S_n$ of part a).

c) If there exists an index $j \in \{1, ..., n\}$ such that $x_j^0 = 0$, but no index i such that $x_i^0 = 1$, then the sets are $S_0, S_1, ..., S_{\beta-1}$ of part a)*, where*

$$\beta := \min\{1 \leq j \leq n | \ x_j^0 = 0\}. \tag{10}$$

d) If there exist an index $i \in \{1, ..., n\}$ such that $x_i^0 = 1$, and an index j such that $x_j^0 = 0$, then the sets are $S_1, ..., S_{\beta-1}$ of part a)*, where β is given by* (10)*.*

Proposition 1 ([18])**.** *For any $x^0 \in \mathcal{B}^n$ the sets $S_i(x^0), 1 \leq i \leq n$, are max-min convex.*

In the following $[a, c]$ denotes the ordinary interval on the real line $\{b\colon a \leq b \leq c\}$, provided $a \leq c$ (and possibly $a = c$).

We investigate the separation of a box $B = [\underline{x}_1, \overline{x}_1] \times \ldots \times [\underline{x}_n, \overline{x}_n]$ from a max-min convex set $C \subseteq \mathcal{B}^n$, by which we mean that there exists a set S described in Definition 1, which contains C and avoids B.

Assume that $\overline{x}_1 \geq \ldots \geq \overline{x}_n$ and suppose that $t(B)$ is the greatest integer such that $\overline{x}_{t(B)} \geq \underline{x}_i$ for all $1 \leq i \leq t(B)$. We will need the following condition:

$$\begin{gathered} \text{If } (\overline{x}_1 = 1) \ \& \ (y_l \geq \underline{x}_l, 1 \leq l \leq n) \ \& \ (\overline{x}_l < y_l \text{ for some } l \leq t(B)), \\ \text{then } y \notin C. \end{gathered} \tag{11}$$

Note that if the box is reduced to a point and if $\overline{x}_1 = 1$, then $\overline{x}_l = 1$ for all $l \leq t(B)$ so that $\overline{x}_l < y_l$ is impossible. So (11) always holds true in the case of a point.

The formulation of our main result will also use an *oracle* answering the question, whether or not a given max-min convex set $C \subseteq \mathcal{B}^n$ lies in a semispace S. As in the conventional convex geometry or tropical convex geometry, this question can be answered in $O(mn)$ time if C is a convex hull of m points. Indeed it suffices to answer whether any of the inequalities defining S is satisfied for each of the m points generating C.

Theorem 1. *Let $B = [\underline{x}_1, \overline{x}_1] \times \ldots \times [\underline{x}_n, \overline{x}_n]$, and let $C \subseteq \mathcal{B}^n$ be a max-min convex set avoiding B. Suppose that B and C satisfy* (11)*. Then there is a set S described by Definition 1, which contains C and avoids B. This set is constructed in no more than $n+1$ calls to the oracle.*

Proof. If $\overline{x}_i < 1$ for all i, then we try to separate B from C by $S_0(\overline{x}_1, \ldots, \overline{x}_n)$ given by (7). Suppose we fail. Then there exists $y \in C$ such that $y_i \leq \overline{x}_i$ for all i.

Otherwise, $1 = \overline{x}_1 \geq \ldots \geq \overline{x}_n$. Let $\underline{x}_l = \max\limits_{k \leq t(B)} \underline{x}_k$, and define $u \in \mathcal{B}^n$ by

$$u_i = \begin{cases} \underline{x}_l, & \text{if } i \leq t(B), \\ \overline{x}_i, & \text{if } i > t(B). \end{cases} \tag{12}$$

It follows from the definition of $t(B)$ that $\underline{x}_l = \max\limits_{1 \leq i \leq n} u_i$.

We try to separate B from C by $S_l(u)$, which is given by (8) or (9). If we fail then there exists $y \in C$ such that $y_i \leq \overline{x}_i$ for all $i > t(B)$ (and trivially $y_i \leq \overline{x}_i$ for $\overline{x}_i = 1$).

Thus we either separate C from B, or there is a point $y \in C$ such that $y_i \leq \overline{x}_i$ for all $i > t(B)$. Condition (11) and $B \cap C = \emptyset$ assure that there is at least one i such that $y_i < \underline{x}_i$. Indeed, otherwise if $\underline{x}_i \leq y_i$ for all i and $y_i \leq \overline{x}_i$ for all $i \leq t(B)$ then $y \in B$; and if $\underline{x}_i \leq y_i$ for all i and $\overline{x}_i < y_i$ for some $i \leq t(B)$ then $y \notin C$ by condition (11).

Now assume without loss of generality that $\underline{x}_1 \geq \ldots \geq \underline{x}_n$ (the order of $\overline{x}_i$ is now arbitrary).

The set $\{1, \ldots, n\}$ is naturally partitioned by the following procedure. See Figure 2 for an illustration. The segments $[\underline{x}_i, \overline{x}_i]$ are drawn vertically and counted from left to right.

Let s_1 be the smallest number such that $\underline{x}_{s_1} \leq \overline{x}_i$ for all $i = s_1, \ldots, n$.



FIGURE 2. The choice of s_n

If $s_1 \neq 1$ then there exists $t_1 \in \{s_1, \ldots, n\}$ such that $\underline{x}_{s_1-1} > \overline{x}_{t_1}$. In this case let T_1 be the set of such t_1. Otherwise if $s_1 = 1$ we take $T_1 = \{1, \ldots, n\}$. In Figure 2 one has $T_1 = \{12\}$.

We define

$$a_1 = \min\{\overline{x}_i \colon i \in T_1\}. \tag{13}$$

We have

$$\underline{x}_i \leq a_1 \leq \overline{x}_i \quad \forall i = s_1, \ldots, n. \tag{14}$$

Thus a_1 is a common level in all intervals $[\underline{x}_{s_1}, \overline{x}_{s_1}], \ldots, [\underline{x}_n, \overline{x}_n]$, but not $[\underline{x}_{s_1-1}, \overline{x}_{s_1-1}]$.

If $s_1 = 1$ then we stop. Otherwise we proceed by induction. Let s_k be the smallest number such that $\underline{x}_{s_k} \leq \overline{x}_i$ for all $i \in \{s_k, \ldots, n\} \backslash T_1 \cup \ldots \cup T_{k-1}$. Note that $s_k < s_{k-1}$. If $s_k \neq 1$ then there exists $t_k \in \{s_k, \ldots, n\}$ such that $\underline{x}_{s_k-1} > \overline{x}_{t_k}$. In this case let T_k be the set of such t_k. Otherwise if $s_k = 1$, then define $T_k := \{1, \ldots, n\} \backslash T_1 \cup \ldots \cup T_{k-1}$.

We take

$$a_k = \min\{\overline{x}_i \colon i \in T_k\}. \tag{15}$$

We have

$$\underline{x}_i \le a_k \le \overline{x}_i \quad \forall i = \{s_k, \dots n\} \backslash T_1 \cup \dots \cup T_{k-1}. \tag{16}$$

Thus a_k is a common level in all intervals $[\underline{x}_{s_k}, \overline{x}_{s_k}], \dots, [\underline{x}_n, \overline{x}_n]$ excluding the intervals with indices in $T_1 \cup \dots \cup T_{k-1}$ which are below that level. The interval $[\underline{x}_{s_k-1}, \overline{x}_{s_k-1}]$ is above a_k (and possibly several other such levels going into T_k).

In Figure 2, the sets T_i are $T_1 = \{12\}, T_2 = \{10\}, T_3 = \{8\}, T_4 = \{7, 9, 11\}, T_5 = \{4\}, T_6 = \{1, 2, 3, 5, 6\}$.

Next we recall our point $y \in C$. It has $y_i < \underline{x}_i$ for some i. Denote $K = \{i \colon y_i > \overline{x}_i\}$. Pick the greatest i such that $y_i < \underline{x}_i$ (note that for such i we necessarily have $\underline{x}_i > 0$), and let $s_k \le i < s_{k-1}$, which implies $\underline{x}_j \le y_j \le \overline{x}_j$ for all $j \in \{s_{k-1}, \dots, n\} \backslash K$. We try to separate B from C by the sets

$$S_i(u^i) = \{x \in \mathcal{B}^n \colon x_i < \underline{x}_i \text{ or } x_j > \overline{x}_j \text{ for some } j \in T_1 \cup \dots \cup T_{k-1}\}, \tag{17}$$

where u^i can be defined by

$$u^i_l = \begin{cases} \underline{x}_l, & l < i, \\ \underline{x}_i, & l \ge i \text{ and } l \notin T_1 \cup \dots \cup T_{k-1}, \\ \overline{x}_l, & l \in T_1 \cup \dots \cup T_{k-1}, \end{cases} \tag{18}$$

for all i with $y_i < \underline{x}_i$ and $s_k \le i < s_{k-1}$. Indeed, (17) is of the form (8) or (9), where u^i is substituted for x^0.

Suppose the separation always fails. Then it gives us points $x^i \in C$ such that

$$x^i_i \ge \underline{x}_i \text{ and } x^i_j \le \overline{x}_j \ \forall j \in T_1 \cup \dots \cup T_{k-1}. \tag{19}$$

Then (16) implies that

$$\underline{x}_i \le a_k \wedge x^i_i \le \overline{x}_i \text{ and } a_k \wedge x^i_j \le \overline{x}_j \ \forall j = s_k, \dots, n, \tag{20}$$

since $a_k \in [\underline{x}_j, \overline{x}_j]$ for $j \in \{s_k, \dots, n\} \backslash T_1 \cup \dots \cup T_{k-1}$ by (16), and we use (19) for $j \in T_1 \cup \dots \cup T_{k-1}$. The point

$$z = \bigoplus_i a_k \wedge x^i \oplus y \in C \tag{21}$$

will be in some sense better than y. Indeed, (20) implies that $\underline{x}_i \leq z_i \leq \overline{x}_i$ for all $i \in \{s_k, \ldots, n\}\backslash K$, versus $\underline{x}_i \leq y_i \leq \overline{x}_i$ for all $i = \{s_{k-1}, \ldots, n\}\backslash K$. As $z \geq y$ we have $z_i > \overline{x}_i$ for all $i \in K$.

Proceeding with this improvement we obtain a point z which satisfies $\underline{x}_i \leq z_i$ for all i and $z_i \leq \overline{x}_i$ for all $i \in \{1, \ldots, n\}\backslash K$. This contradicts either $B \cap C = \emptyset$, or condition (11). This contradiction shows that we should succeed with separation at some stage. Clearly, the number of calls to the oracle does not exceed $n + 1$. □

We note that Theorem 1 also yields a method which verifies condition (11) in no more than $n + 1$ calls to the oracle.

The box B can be a point and in this case condition (11) always holds true. Therefore, some known results on max-min semispaces [18] can be deduced from Theorem 1. The following statement is an immediate corollary of Theorem 1 and Proposition 1.

Corollary 1 ([18])**.** *Let $x \in \mathcal{B}^n$ and $C \subseteq \mathcal{B}^n$ be a max-min convex set avoiding x. Then C is contained in one $S_i(x), 1 \leq i \leq n$, as in Definition 1. Consequently these sets are the family of semispaces at x.*

Proof. The proof of Theorem 1 applied to $B = \{x\}$ shows that any max-min convex set avoiding x is contained in one of the sets $S_i(x)$. Proposition 1 implies that these sets are max-min convex and do not contain x. Obviously, they are not included in each other. If $S_i(x)$ is not maximal, let S be a max-min convex set strictly containing $S_i(x)$. Then Theorem 1 implies that there exists other $S_j(x), i \neq j$, such that $S \subset S_j(x)$. But this implies $S_i(x) \subset S_j(x)$, a contradiction. Hence $S_i(x)$ are all maximal and $\{S_i(x)\}_i$ is the family of semispaces at x. □

Thus we recover a result of [18] that Definition 1 actually yields all semispaces at a given point.

We now show that separation by semispaces is impossible when B and C do not satisfy (11).

Theorem 2. *Suppose that* $B = [\underline{x}_1, \overline{x}_1] \times \ldots \times [\underline{x}_n, \overline{x}_n]$ *and max-min convex set* $C \subseteq \mathcal{B}^n$ *are such that* $B \cap C = \emptyset$ *but the condition* (11) *does not hold. Then there is no semispace that contains* C *and avoids* B*.*

Proof. We assume that $\overline{x}_1 \geq \ldots \geq \overline{x}_n$. Since (11) does not hold, we have $\overline{x}_1 = 1$. Also there exists $z \in C$, such that for some indices $k \leq t(B)$ we have

$$\max_i \{\underline{x}_i \colon i \leq k\} \leq \overline{x}_k < z_k, \tag{22}$$

but

$$\underline{x}_i \leq z_i \leq \overline{x}_i \tag{23}$$

for all the others. Any semispace of the type S_0 given by (7) intersects with B since $\overline{x}_1 = 1$. If we assume by contradiction that a separating semispace exists, then it must be of the type $S_i(x^0)$ given by (8) or (9). Further, we claim that this semispace must contain the set $\{x \colon x_k > x_k^0\}$ for some k such that $z_k > x_k^0 \geq \overline{x}_k$, otherwise it either does not contain z or it intersects with B. Indeed $z \in S_i(x^0)$ implies

$$z_i < x_i^0, \text{ or } z_j > x_j^0 \text{ for } j \text{ such that } x_i^0 > x_j^0. \tag{24}$$

If $z_j > x_j^0$ is true for some j such that $x_j^0 < \overline{x}_j$, then $y \in S_i(x^0)$ for each $y \in B$ with $y_j = \overline{x}_j$, hence $B \cap S_i(x^0) \neq \emptyset$, a contradiction. If $z_i < x_i^0$ is true, then $x_i^0 > \underline{x}_i$ implying $y \in S_i(x^0)$ for each $y \in B$ with $y_i = \underline{x}_i$, hence again $B \cap S_i(x^0) \neq \emptyset$, a contradiction. Thus $z_k > x_k^0$ must hold true for at least one k, and necessarily with $x_k^0 \geq \overline{x}_k$.

This also implies that $i \neq k$, for the type of the semispace above. Then we must have $x_i^0 > x_k^0 \geq \overline{x}_k$ and $\{x \mid x_i < x_i^0\} \subseteq S_i(x^0)$. If $i < k$ then $x_i^0 > \overline{x}_k \geq \underline{x}_i$ due to (22), and if $i > k$ then also $x_i^0 > \overline{x}_k \geq \overline{x}_i \geq \underline{x}_i$ by the ordering of $\overline{x}_i$. Hence the set $\{x \colon x_i < x_i^0\}$, which is contained in $S_i(x^0)$, intersects with B and the separation is impossible. □

Remark 1. *A simple example of interval non-separation is shown in Figure 2. The box is* $B = [0, 1] \times [a, b]$ *where* $0 \leq a \leq b < 1$ *and the convex set is* $C = \{z\}$ *where* $z = (z_1, z_2)$ *with* $z_2 > b$*. Note that* B *and* C *do not satisfy condition* (11)*.*



FIGURE 3. Forbidden 2-dim interval separation of convex sets

Remark 2. *Theorem 1 can be easily modified to allow any case, if in addition to Definition 1 we also allow the sets*

$$S_0^M(x^0) = \{x\colon x_i > x_i^0 \text{ for some } i \in M\}. \tag{25}$$

By Corollary 1 these sets cannot be semispaces. They are hemispaces *in the sense that both the set and its complement are max-min convex. The condition $C \subseteq S_0^M(x^0)$ can be verified by the same type of oracle as in Theorem 1.*

3. Separation of two max-min convex sets

In this section we investigate the separation of two disjoint closed max-min convex sets by a box and by a box and a semispace.

We recall the structure of 2-dimensional max-min segments as presented in [17]. Pictures of all types of max-min segments are shown in Figure 4, taken from [17].

Theorem 3. *Let $C_1, C_2 \in \mathcal{B}^2$, $C_1 \cap C_2 = \emptyset$, be two closed max-min convex sets. Then there exist a permutation $i : \{1,2\} \to \{1,2\}$ and a box $B \subset \mathcal{B}^2$ such that $C_{i(1)} \subset B$ and $B \cap C_{i(2)} = \emptyset$.*

Proof. Let

$$\begin{aligned} x_c &:= \max\{x | (x,y) \in C_1 \text{ for some } y\}, \\ y_c &:= \max\{y | (x,y) \in C_1 \text{ for some } x\}. \end{aligned} \tag{26}$$



FIGURE 4. 2-dim max-min segments

As C_1 is compact, there exist $(x_c, y), (x, y_c) \in C_1$, and the convexity of C_1 implies that

$$c := (x_c, y_c) = (x_c, y) \oplus (x, y_c) \in C_1. \tag{27}$$

Let

$$\begin{aligned} x_a &:= \min\{x | (x, y) \in C_1 \text{ for some } y\}, \\ y_b &:= \min\{y | (x, y) \in C_1 \text{ for some } x\}. \end{aligned} \tag{28}$$

Consider the points in C_1, guaranteed again by compactness:

$$\begin{aligned} a &:= (x_a, y_a), \\ b &:= (x_b, y_b). \end{aligned} \tag{29}$$

The values y_a and x_b are chosen arbitrarily.

The smallest box in $\mathcal{B}^2$ containing the convex set C_1 is $B_0 := [x_a, x_c] \times [y_b, y_c]$. The point c is the upper right corner of B_0.



FIGURE 5. 2-dim box separation

We need the following Lemma, which can be proved by drawing all possible special cases and using the structure of max-min segments shown on Figure 4. This proof is routine and will be omitted.

Lemma 1. *The box B_0 can be partitioned as $B_0 = T_0 \cup T_1 \cup T_2 \cup T_3$, where*

$$T_0 = \{\alpha \wedge a \oplus \beta \wedge b \oplus \gamma \wedge c \colon \alpha \oplus \beta \oplus \gamma = 1\},$$

$$T_1 = B_0 \cap \{(x, y) \colon x < x_b,\ y < y_a\},$$

$$T_2 = B_0 \cap \{(x, y) \colon y > y_a,\ x < x_c,\ y > x\},$$

$$T_3 = B_0 \cap \{(x, y) \colon x > x_b,\ y < y_c,\ y < x\}.$$

All regions T_0, T_1, T_2, T_3 are max-min convex (or possibly empty).

The regions T_0, T_1, T_2, T_3 are shown in Figure 5.

Evidently $T_0 \subseteq C_1$ (note that T_0 is the max-min convex hull of a, b, c). In particular, the max-min segments $[a, b]_M, [a, c]_M, [b, c]_M$ are included in C_1 and any point from C_2 stays away from them. The other regions may contain points from both C_1 and C_2.

We show that if the convex set C_2 intersects with one of the regions T_1 and T_2, then there is a box $B_1 \subset \mathcal{B}^2$ such that $C_2 \subset B_1$ and $B_1 \cap C_1 = \emptyset$. Due to the symmetry about the main diagonal, there is no need to consider the case where C_2 intersects with T_3.

Case 1. Assume C_2 intersects with the region T_1.

The intersection C_2' of C_2 with region T_1 is max-min convex (as the intersection of two max-min convex sets). Thus there exists a point $(x_M, y_M) \in C_2'$, away from the boundary of C_1, and consequently away from the segment $[a,b]_M$, that has the maximum x-coordinate and maximum y-coordinate for C_2'.

We show that C_2 is included in the box $B_1 = [0, x_M] \times [0, y_M]$.

Assume by contradiction that there exists $(x', y') \in C_2$ such that $x' > x_M$ or $y' > y_M$, then $(x'', y'') := (x_M, y_M) \oplus (x', y')$ has either $x'' = x_M$ and $y'' = y' > y_M$, or $x'' = x' > x_M$ and $y'' = y_M$, or $(x'', y'') = (x', y')$. If $x'' < x_b$ and $y'' < y_a$ then $(x'', y'') \in C_2'$, which contradicts the maximality of x_M and y_M. Otherwise, the segment $[(x_M, y_M), (x'', y'')]_M$ intersects with $[a,b]_M$ and hence $C_1 \cap C_2 \neq \emptyset$, a contradiction.

We show that the box B_1 does not intersect with C_1. Assume that there exists $(x,y) \in B_1 \cap C_1$. Then $x \leq x_M$ and $y \leq y_M$. There exist $(x', y_M) \in [a,b]_M$ and $(x_M, y') \in [a,b]_M$ such that $x' > x_M$ and $y' > y_M$. Using these points, we obtain that

$$\begin{aligned} (x_M, y_M) &= (x,y) \oplus x_M \wedge (x', y_M), \quad \text{if } x_M \geq y_M \\ (x_M, y_M) &= (x,y) \oplus y_M \wedge (x_M, y'), \quad \text{if } x_M \leq y_M. \end{aligned} \tag{30}$$

In both cases $(x_M, y_M) \in C_1$ and hence $C_1 \cap C_2 \neq \emptyset$, a contradiction.

Case 2. Assume now that C_2 intersects with T_2, and let $C_2' := C_2 \cap T_2$. Let x_M be the largest x coordinate of a point in C_2' and y_M the lowest y-coordinate of a point in C_2'. Let $(x_0, y_M), (x_M, y_0) \in C_2'$. From the definition of T_2 we have $x_0 \leq y_M$ and $x_M \leq y_0$. Let $[t_1, t_2] := [x_a, x_c] \cap [y_a, y_c]$ (where all segments are ordinary on the real line).

If $y_M \leq x_M$, then due to convexity

$$\begin{aligned} (x_0, x_0) &= (x_0, y_M) \oplus x_0 \wedge (x_M, y_0) \in C_2', \\ (y_M, y_M) &= (x_0, y_M) \oplus y_M \wedge (x_M, y_0) \in C_2', \end{aligned} \tag{31}$$

and hence the whole diagonal (and max-min) segment $[(x_0, x_0), (y_M, y_M)]_M$ is included in C_2'. It can be observed that any point in the closure of T_2 that belongs to the main diagonal lies in $[a,c]_M$ which is in C_1. Thus $C_1 \cap C_2 \neq \emptyset$, a contradiction, hence we must have $y_M > x_M$.

When $y_M > x_M$, due to convexity we have

$$(x_M, y_M) = (x_0, y_M) \oplus y_M \wedge (x_M, y_0) \in C_2'. \tag{32}$$

In this case we claim that C_2 is contained in the box $B_1 := [0, x_M] \times [y_M, 1]$, which avoids C_1.

Assume by contradiction that there exists $(x', y') \in C_2$ which does not lie in B_1. This implies that $x' > x_M$ or $y' < y_M$. We also have $y_M > x'$ and $y' > x_M$, otherwise the segment $[(x', y'), (x_M, y_M)]_M$ has points on the main diagonal, in which case it intersects with $[a, c]_M$. Consider the combinations

$$\begin{aligned} &(x', y_M) = y_M \wedge (x', y') \oplus (x_M, y_M), \quad \text{if } x' > x_M, \\ &(x_M, y') = (x', y') \oplus y' \wedge (x_M, y_M), \quad \text{if } y' < y_M \text{ and } x' \leq x_M. \end{aligned} \tag{33}$$

Thus we obtain either $(x', y_M) \in C_2$ with $x' > x_M$, or $(x_M, y') \in C_2$ with $y' < y_M$ and $x' \leq x_M$, leading to a contradiction with the maximality of x_M or the minimality of y_M.

To prove that B_1 avoids C_1, assume by contradiction that there exists $(x, y) \in C_1$ where $x \leq x_M$ and $y \geq y_M$. We observe that there is a point $(x_M, y') \in [a, c]_M$, where $y' \leq y_M$. Using this point we obtain

$$(x_M, y_M) = y_M \wedge (x, y) \oplus (x_M, y'), \tag{34}$$

which implies $(x_M, y_M) \in C_1$, hence $C_1 \cap C_2 \neq \emptyset$, a contradiction. □

Theorem 4. *Let $C_1, C_2 \in \mathcal{B}^2$, $C_1 \cap C_2 = \emptyset$, be two closed max-min convex sets that are away from the boundary of $\mathcal{B}^2$. Then there exist a permutation $i : \{1, 2\} \to \{1, 2\}$, a box $B \subset \mathcal{B}^2$ and a semispace $S \subset \mathcal{B}^2$ such that $C_{i(1)} \subset B, C_{i(2)} \subset S$ and $B \cap S = \emptyset$.*

Proof. The statement follows from Theorem 1 and Theorem 3. Indeed, Theorem 3 implies that either the minimal containing box of C_1 does not intersect with C_2, or the minimal containing box of C_2 does not intersect with C_1. The condition (11) is satisfied due to the fact that the convex sets are away from the boundary of $\mathcal{B}^n$ and hence so are the minimal containing boxes. Applying Theorem 1 we obtain the statement. □



FIGURE 6. Forbidden 3-dim box-semispace separation

Remark 3. *We observe that Theorem 3 and Theorem 4 are not valid in dimension 3 or higher.*

Let C_1 *be the max-min segment* $[(a,a,a),(b,b,b)]_M$ *and* C_2 *be the max-min segment* $[(b,a,a),(a,b,a)]_M$*, where* $0 \leq a < b \leq 1$*. It follows from* [17] *that* C_1 *is part of the main diagonal, and that* C_2 *is the concatenation of two pieces with parametrizations* $\{(t,b,a)|a \leq t \leq b\}$ *and* $\{(b,t,a)|a \leq t \leq b\}$*. It follows from Figure 6 that the smallest box containing* C_1 *is* $[a,b]^3$ *and the smallest box containing* C_2 *is* $[a,b]^2 \times \{a\}$*. Since one box is completely included in the other, box separation or box-semispace separation of* C_1 *and* C_2 *is not possible.*

REFERENCES

[1] X. Allamegeon, S. Gaubert, and R. Katz. The number of extreme points of tropical polyhedra. E-print arXiv:math/0906.3492v1, 2009.

[2] G. Birkhoff. *Lattice theory.* American Mathematical Society, Providence, 1993.

[3] K. Cechlárová. Eigenvectors in bottleneck algebra. *Linear Algebra Appl.*, 175:63–73, 1992.

[4] G. Cohen, S. Gaubert, J.P. Quadrat, and I. Singer. Max-plus convex sets and functions. In G. Litvinov and V. Maslov, editors, *Idempotent Mathematics and Mathematical Physics*, volume 377 of *Contemporary Mathematics*, pages 105–129. AMS, Providence, 2005. E-print arXiv:math/0308166.

[5] M. Develin and B. Sturmfels. Tropical convexity. *Documenta Math.*, 9:1–27, 2004. E-print arXiv:math/0308254.

[6] M. Fiedler, J. Ramík, J. Nedoma, J. Rohn, and K. Zimmermann. *Linear optimization problems with inexact data*. Springer Verlag, 2006.

[7] S. Gaubert and R. Katz. *Max-plus convex geometry*, volume 4136 of *Lecture Notes in Computer Sciences*, pages 192–206. Springer, New York, 2006.

[8] S. Gaubert and S. Sergeev. Cyclic projectors and separation theorems in idempotent convex geometry. *Journal of Math. Sci.*, 155(6):815–829, 2008. E-print arXiv:math/0706.3347.

[9] M. Gavalec. *Periodicity in Extremal Algebra*. Gaudeamus, Hradec Králové, 2004.

[10] J. Golan. *Semirings and their applications*. Kluwer, Dordrecht, 2000.

[11] G.L. Litvinov, V.P. Maslov, and G.B. Shpiz. Idempotent functional analysis. algebraic approach. *Math. Notes (Moscow)*, 69(5):758–797, 2001.

[12] G.L. Litvinov and A.N. Sobolevskiĭ. Idempotent interval analysis and optimization problems. *Reliable Computing*, 7(5):353–377, 2001.

[13] V. Nitica. The structure of max-min hyperplanes. Submitted to Linear Algebra Appl., 2009.

[14] V. Nitica and S. Sergeev. On hyperplanes and semispaces in max-min convex geometry. Submitted to Kybernetika, 2009.

[15] V. Nitica and I. Singer. Max-plus convex sets and max-plus semispaces. i. *Optimization*, 56:171–205, 2007.

[16] V. Nitica and I. Singer. Max-plus convex sets and max-plus semispaces. ii. *Optimization*, 56:293–303, 2007.

[17] V. Nitica and I. Singer. Contributions to max-min convex geometry. i: Segments. *Linear Algebra Appl.*, 428(7):1439–1459, 2008.

[18] V. Nitica and I. Singer. Contributions to max-min convex geometry. ii: Semispaces and convex sets. *Linear Algebra Appl.*, 428(8-9):2085–2115, 2008.

[19] B. Semančíková. Orbits in max-min algebra. *Linear Algebra Appl.*, 414:38–63, 2006.

[20] S. N. Sergeev. Algorithmic complexity of a problem of idempotent convex geometry. *Math. Notes (Moscow)*, 74(6):848–852, 2003.

[21] I. Singer. *Abstract convex analysis*. Wiley-Interscience, 1997.

[22] K. Zimmermann. A general separation theorem in extremal algebras. *Ekonom.-Mat. Obzor (Prague)*, 13:179–201, 1977.

[23] K. Zimmermann. Convexity in semimodules. *Ekonom.-Mat. Obzor (Prague)*, 17:199–213, 1981.

Department of Mathematics, West Chester University, PA 19383, USA, and Institute of Mathematics, P.O. Box 1-764, Bucharest, Romania

E-mail address: vnitica@wcupa.edu

University of Birmingham, School of Mathematics, Watson Building, Edgbaston B15 2TT, UK

E-mail address: sergiej@gmail.com